

# Onslaught®

## MICROENCAPSULATED INSECTICIDE

**ACTIVE INGREDIENT:**

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate . . . . . 6.40%

\*OTHER INGREDIENTS . . . . . 93.60%  
100.00%

\*Contains petroleum distillates

### KEEP OUT OF REACH OF CHILDREN CAUTION

**FIRST AID**

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Do not give any liquid to the person. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**NOTE TO PHYSICIAN:** Contains petroleum distillate - vomiting may cause aspiration pneumonia.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with

soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage system. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Remove pets and birds and cover fish aquariums before spraying.

See Back Panel for Additional Precautionary Language

Manufactured by:




8810 Tenth Avenue North • Minneapolis, MN 55427

EPA Reg. No. 1021-1815    EPA Est. No. 1021-MN-2<sup>A</sup>, 63416-MN-1<sup>B</sup>,  
Rev. 0613-0214    70815-GA-1<sup>C</sup>, 39578-TX-1<sup>D</sup>,  
F28371-107    44616-MO-1<sup>E</sup>, 47000-IA-1<sup>F</sup>

Superscript is the first character of the lot number.

**NET CONTENTS: 1.0 GALLON**

PULL HERE TO OPEN / PRESS TO RESEAL



**FIRST AID**

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give <b>any</b> liquid to the person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or a doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN:</b> Contains petroleum distillate – vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wear protective eyewear (goggles, face shield or safety glasses with side shields). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish and other aquatic invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage system. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirement of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Remove pets and birds and cover fish aquariums before spraying.

**For control of insects, indoors and outdoors, in food and non-food areas such as, but not limited to:** Homes, Schools, Warehouses, Office Buildings, Apartment Buildings, Theatres, Hotels, Industrial Buildings, Motels, Kennels, Livestock Housing, Food Processing Plants, Food Service Establishments, Restaurants, Supermarkets and Grocery Stores, Transportation Equipment, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments. Also for use on Backyards, Lawns, Trees, Ornamental Landscaping, Recreational Areas, Parks and Athletic Fields.

**Insects Controlled:**

Ants  
Aphids  
Asian Lady Beetles  
Bed Bugs  
Beetles  
Box Elder Bugs  
Brown Marmorated Stink Bugs  
Carpet Beetles  
Centipedes

Cockroaches  
Crickets  
Dog Ticks  
Earwigs  
Fire Ants  
Firebrats  
Fleas  
Gnats  
Hornets

Kudzu Bugs  
Lice  
Mealy Bugs  
Midges  
Millipedes  
Mites  
Mosquitoes  
Pillbugs  
Psocids

Scales  
Scorpions  
Silverfish  
Spiders  
Springtails  
Swarming Termites  
Ticks  
Wasps  
Waterbugs

**Flying Insects Including:**

Blow Flies  
Chiggers  
Clothes Moths  
Cluster Flies

Crane Flies  
Deer Flies  
Face Flies  
Fruit Flies

Fungus Gnats  
Gnats  
Horn Flies  
Houseflies

Mosquitoes  
Small Flying Moths  
Whiteflies

**Biting and Stinging Pests:**

Bed Bugs  
Bees  
Biting Flies  
Chiggers  
Deer Flies

Deer Ticks  
Dog Ticks  
Fire Ants  
Fleas  
Gnats

Hornets  
Lice  
Mosquitoes  
Scorpions

Spiders  
Ticks  
Wasps  
Yellow Jackets

**Stored Product Pests including:**

Angoumois Grain Moths  
Ants  
Cadelles  
Cigarette Beetles  
Confused Flour Beetles  
Dark Mealworms  
Dried Fruit Beetles  
Drugstore Beetles

Flat Grain Beetles  
Fruit Flies  
Grain Mites  
Grain Moths  
Granary Weevils  
Indian Meal Moths  
Lesser Grain Borers  
Maize Weevils

Meal Moth Larvae  
Mediterranean Flour Moths  
Merchant Grain Beetles  
Red Flour Beetles  
Rice Weevils  
Rusty Grain Beetles  
Saw-toothed Grain Beetles

Skippers  
Spider Beetles  
Spider Mites  
Square-necked Grain Beetles  
Tobacco Moths  
Yellow Mealworms

**Livestock Premise Pests:**

Bed Bugs  
Carrion Beetles  
Chiggers  
Darkling Beetles  
(lesser mealworm)

Deer Flies  
Face Flies  
Fleas  
Flies

Hide Beetles  
Horn Flies  
Horse Flies  
Lice

Litter Beetles  
Mites  
Mosquitoes  
Stable Flies

**Wood-Destroying Pests including:**

Carpenter Ants  
Carpenter Bees

Deathwatch Beetles  
Furniture Beetles

Old House Borers  
Powder Post Beetles

Round-headed House Borers  
Swarming Termites

**Plant Pests including:**

Alfalfa Caterpillars	Cranberry Weevils	Leaf Miners	Pear Slugs	Sap Beetles
American Plum Borers	Cucumber Beetles	Leaf Rollers	Pecan Leaf Phylloxera	Scales
Annual Bluegrass Weevils	Cutworms	Leaf Tiers	Pecan Nut Casebearers	Sod Webworms
Aphids	Diamondback Moths	Leafhoppers	Pecan Spittlebugs	Sowbugs
Apple Maggots	Elm Leaf Beetles	Lesser Appleworms	Pecan Stem Phylloxera	Spiders
Armyworms	European Pine Sawflies	Lesser Peach Tree Borers	Pecan Weevils	Spittlebugs
Artichoke Plume Moth	Fall Webworms	Loopers	Pepper Weevils	Stink Bugs
Bagworms	Filbert Worms	Lygus Bugs	Periodical Cicadas	Tarnished Plant Bugs
Balsam Woolly Adelgids	Flea Beetles	Mexican Bean Beetles	Pickleworms	Tent Caterpillars
Beet Armyworms	Grasshoppers	Mites	Pillbugs	Thrips
Billbugs	Green Cloverworms	Mole Crickets	Pine Chafers	Tobacco Hornworms
Blueberry Spanworms	Green Fruitworms	Naval Orangeworms	Pine Coreid Bugs	Tufted Apple Bud Moths
Brown Marmorated Stink Bugs	Gypsy Moth Caterpillars	Northern Pine Weevils	Plant Bugs	Velvet Bean Caterpillars
Cherry Fruit Flies	Hickory Shuckworms	Oriental Fruit Moths	Plum Curculios	Walnut Husk Flies
Cherry Fruitworms	Imported Cabbageworms	Painted Lady Caterpillars	Red Pine Sawflies	Western Bean Cutworms
Chinch Bugs	Japanese Beetles	Pea Weevils	Red-headed Pine Sawflies	and Whiteflies
Codling Moths	Kudzu Bugs	Peach Tree Borers	Red-striped Fireworms	
Cowpea Curculios	Lace Bugs	Peach Twig Borers	Rindworms	
Cranberry Fruitworms	Leaf Feeding Caterpillars	Pear Psyllid	Salt Marsh Caterpillars	

**GENERAL INFORMATION**

Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, schools, sports facilities, etc.) in the immediate area where occupants are present.

ONSLAUGHT® Microencapsulated Insecticide is a microencapsulated suspension concentrate containing 0.54 pound active ingredient per gallon.

ONSLAUGHT® Microencapsulated Insecticide is designed to provide long residual activity against pests listed on this label when applied as surface or spot treatments, injected into wall voids, or as a crack and crevice spray in and around residential and commercial structures, as well as on turf and landscaping. ONSLAUGHT® Microencapsulated Insecticide may be sprayed on any surface that will not be stained by water. DO NOT use this product as a fogger or apply it as a space spray. Do not apply in electrical conduits, junction and switch boxes, motor housings, or other electrical equipment due to shock hazard from water-based spray.

Use only in areas described. Remove pets and cover fish aquariums and pets' food and water dishes before spraying. Keep all people (especially children) and pets out of areas being treated and restrict access to these areas until all surfaces are dry. DO NOT leave product where children or pets can come into contact with it. DO NOT allow spray to contact food or food-containing surfaces, feed, or water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated with this product. Food utensils such as teaspoons and measuring cups must not be used for food purposes after use with pesticides.

**APPLICATION IN FOOD PROCESSING/HANDLING ESTABLISHMENTS:**

Application is Permitted Within Food and Non-Food Areas of Food Service and Handling Establishments (Places Other than Private Residences) Including But Not Limited to: Restaurants, Meat Processing Plants, Grocery Stores, Bakeries, Food Manufacturing and Processing Establishments, and Food Warehousing Establishments. Do not treat establishments where livestock feed is present. Food areas include areas for receiving, storing, packing, canning, bottling, wrapping, boxing, preparing, edible waste storage, and enclosed processing systems, mills, and dairies. Serving areas are places where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, garages, mop closets, and storage after canning or bottling.

**General Surface Applications:** Do not apply ONSLAUGHT® Microencapsulated Insecticide as a general surface application in food areas of the facility when the facility is in operation or food is exposed. Do not apply directly to food. Cover or remove all food and food processing equipment prior to application. After spraying in meat packing plants, bakeries, and other food processing plants, wash with an effective cleaning compound, and then rinse all equipment, benches, shelving, etc. where exposed food will be handled with potable water. Repeat applications as needed, but do not exceed more than one (1) application every (14) days.

**Spot or Crack & Crevice Applications:** Spot or crack and crevice applications may be made while facility is in operation, provided exposed food is covered or removed from the area being treated. Do not apply directly to food or food-handling surfaces. Repeat applications as needed, but do not exceed more than one (1) application every (14) days. Do not apply when food-processing facility is in operation or foods are exposed. Do not apply this spray to surfaces or utensils that may come in contact with food.

In the home, cover exposed food and do not allow spray to contact food surfaces. If spray does contact these surfaces, clean surfaces with soap and water.

## MIXING INSTRUCTIONS

ONSLAUGHT® Microencapsulated Insecticide should be mixed with water and applied with hand pressurized or power operated sprayers. Shake container of ONSLAUGHT® Microencapsulated Insecticide before diluting. Clean spray equipment before use. For dilution, add approximately half the required water to spray tank and then add the appropriate amount of ONSLAUGHT® Microencapsulated Insecticide. Agitate, and slowly add the remaining water. Agitate spray thoroughly before using and also occasionally during use to ensure dispersion. If spray filter screens are used, they should be 50 mesh or larger. Use 0.5 fluid ounce of ONSLAUGHT® Microencapsulated Insecticide in 1.0 gallon of water to make a spray mix containing 0.025% active ingredient. Use 1.0 fluid ounce per gallon of water to make a spray containing 0.05% active ingredient. Use 0.025% solution for light infestations or as a maintenance control rate. Use 0.05% for heavy infestations or as an initial clean-out rate. Apply two-second bursts of spray per square foot of area being treated. Avoid excessive application. Dampen surfaces but not to the point of saturation or runoff. Only dilute ONSLAUGHT® Microencapsulated Insecticide with water.

**Formula for determining the active ingredient content of the finished spray mixture:** The following formula may be used to determine the percent active ingredient that is in the sprayer tank after mixing ONSLAUGHT® Microencapsulated Insecticide:

$$\frac{(6.4) \times (\text{number of fl. oz. of ONSLAUGHT}^{\circledR} \text{ Microencapsulated Insecticide added to tank})}{(\text{Gallons of finished spray mix}) \times (128)} = \begin{array}{l} \text{Percent Active Ingredient} \\ \text{of Spray Mix} \end{array}$$

### Tank Mixing:

ONSLAUGHT® Microencapsulated Insecticide may be tank mixed with an insect growth regulator such as NyGuard® IGR Concentrate or pyrethrum-containing products or any other currently registered pesticides unless expressly prohibited by the product label. The resulting tank mix may be applied in areas where these products are allowed to be sprayed. Do not tank mix ONSLAUGHT® Microencapsulated Insecticide with products containing dichlorvos (DDVP).

### INDOOR USE:

#### Application Rates for Indoor Structural Pests:

Pests	Light Infestation		Heavy Infestation		Application Rate of Mixed Solution	Application Methods
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water		
Crawling Insects	0.025%	0.5 fl oz /gallon	0.050%	1.0 fl oz/ gallon	1 gallon/1000 sq. ft.	Apply as a broadcast surface spray, crack & crevice treatment, spot treatment, or injection into wall voids.
Flying Insects						

#### Crawling Insect Pests Indoors:

Apply as a coarse, wet spray to surfaces where these pests are normally found. Treat floors, baseboards, around doors and windows, in attics, crawl spaces, eaves, corners, closets, walls, utility pipes, storage areas, and all cracks and crevices. Treat underneath sinks, dishwashers, refrigerators, stoves, the underside of shelves, drawers, cabinets, areas behind pipes, and in all places where these insects shelter. Contact as many insects as possible with direct spray. Repeat applications as needed, but do not exceed more than one (1) application every seven (7) days.

#### For Ant Control Indoors:

Apply to ant trails, around garbage receptacles, and near food sources. Also apply around doors and windows and wherever these pests may find entrance to the structure. Remove sources of food through basic hygiene practices whenever possible.

#### Flying Insect Pests Indoors:

Treat insect resting areas such as walls and ceilings, screens, around windows, doors, and light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season, and repeat treatments during periods of heavy infestation.

#### Stinging Insect Pests Indoors:

Apply spray to nests late in the evening when stinging insects are at rest. Thoroughly spray nest, nest entrance, and surrounding areas where insects land or walk. Treat around doors and windows, in attics, crawl spaces, and possible harborage sites or points of entry.

#### For Control of Brown Dog Ticks:

Thoroughly apply as a spot treatment to infested areas such as pet beds and resting areas, nearby cracks and crevices, along and behind baseboards, window and door frames, and localized areas of floor and floor coverings where these pests may be present. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

**For Control of Fleas:**

Thoroughly apply as a fine particle broadcast spray to infested rugs, carpets, and pet resting areas. Prior to treatment, aquariums and fish bowls should be covered, and pet animals should be removed from the area being treated. Do not permit humans or pets to contact treated surfaces until spray has dried. Old pet bedding should be removed and replaced with clean, fresh bedding after treatment. DO NOT SPRAY PETS WITH THIS PRODUCT. Treat dogs and cats with a product registered for use on animals.

**To Control Bed Bugs:**

Thoroughly clean and sanitize mattresses and box springs. Treat mattresses and box springs with an approved pesticide, such as a pyrethrin aerosol. Apply ONSLAUGHT® Microencapsulated Insecticide as a spot treatment to potential harborage sites and migration paths, and cracks and crevices, around baseboards, floorboards, headboards, and walls.

**For Control of Stored Product Pests:**

Spray thoroughly around and into floor drains, non-food conveyors, benches, pipes, pallets, moist areas, storage racks, pieces of equipment, and other areas where stored product pests may be found. Tank-mix or sequential use of an insect growth regulator, such as NyGuard® IGR Concentrate, is recommended to break the insect reproduction cycle. Do not apply this spray to surfaces or utensils that may come in contact with food, since excessive residues in food may result.

**To Treat Voids in Equipment and Structures:**

To kill insects harboring in wall voids and other inaccessible spaces in equipment and structures, use injection equipment designed for deep void applications. Follow the injection equipment manufacturer's recommendations for proper set up and air pressures. Place the applicator tip at or into the void space to be treated. For inaccessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in short bursts, allowing air pressure to push insecticide deep into the space. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry. Do not reapply more than every fourteen (14) days.

**To Kill the Accessible Stages of Listed Granary Insects:** Use this concentrate to treat grain storage facilities, and other listed areas, for stored product pest control. For control of exposed adult and immature stages of stored product pests, apply to cracks, crevices, and other surfaces where the pests have been seen or have harborage. Treat areas where products are stored before filling with the product. Apply at the rates listed above, using one gallon of spray mix per 1,000 square feet of surface area to be treated. Cleaning all areas prior to use will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

Repeat application as needed, but do not exceed more than one application every fourteen (14) days.

**USE IN AND AROUND LIVESTOCK HOUSING:**

For use in and around unoccupied areas of livestock facilities, such as, but not limited to: Barns, Cow and Calf Pens and Hutches, Dairy Barns and Milk Rooms, Hog Barns, Horse Barns, Sheep Barns, Poultry Houses, and Rabbit Hutches. Do not contaminate milk, food, or drinking water. Remove animals from area being treated. Cover feeders and waterers. Do not apply or allow insecticide to drift onto animals. Do not allow animals to enter treated areas until spray solution has dried.

Pests	Light Infestation		Heavy Infestation		Application Methods
	% Active Ingredient	Dilution Rate in Water	% Active Ingredient	Dilution Rate in Water	
Crawling Insects *Including Litter Beetles	0.025%	0.5 fl oz /gallon or sufficient water to cover 1000 sq ft	0.050%	1.0 fl oz /gallon or sufficient water to cover 1000 sq ft	Apply as a general surface spray, crack & crevice treatment, spot treatment, or injection into wall voids.
Flying Insects					

**Crawling Insect Pests in Livestock and Poultry Premises:**

Apply as a general surface, spot treatment, and/or crack and crevice treatment. Apply to floors and vertical and overhead surfaces where crawling insects are or may be present. Treat stanchions, pipes, windows, doors, posts, cage framing, gates, under (but not in) feeders, and other areas where insects hide or congregate. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Litter Beetle Control in Livestock and Poultry Premises:**

To control litter beetles (darkling, hide, and carrion beetles), apply ONSLAUGHT® Microencapsulated Insecticide to walls and floors at cleanout and before reintroduction of animals. Treat areas where beetles frequently occur, such as walls, supports, cages, cage framing, stalls, and around feeders. To reduce immigration of insects, make a perimeter treatment around the outside of building foundations. Apply in a uniform band 1-3 feet up the exterior foundation wall and 3-6 feet out from the foundation.

**Flying Insect Pests in Livestock and Poultry Premises:**

For residual control of flying insects, treat insect resting areas such walls, ceilings, screens, around windows, doors, light fixtures, and other surfaces that attract flying insects. Initiate treatments at the beginning of fly season and repeat treatments during periods of heavy infestation.

**OUTDOOR USE:**

Do not spray in or near fish ponds or other bodies of water.

Do not water the treated area to the point of runoff.

Do not make applications during the rain.

Do not apply directly to sewers or drains, or any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur, except as directed by this label.

Not for use on plants being grown for sale or other commercial use. Not for use in nurseries, sod farms or golf courses.

All outdoor applications, if permitted elsewhere on this label must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses, if allowed elsewhere on this label:

1. Applications to soil or vegetation, as listed on this label;
2. Applications to lawns, turf, and other vegetation, as listed on this label;
3. Applications to the side of a building, up to a maximum height of 3 feet above grade;
4. Application to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
5. Applications around potential pest entry points into buildings, when limited to a surface band not to exceed one inch in width;
6. Applications made through the use of a coarse, low pressure spray to only those portions of surfaces that are directly above bare soil, lawn, turf, mulch or other vegetation, as listed on this label, and not over an impervious surface, drainage or other condition that could result in runoff into storm drains, drainage ditches, gutters, or surface waters in order to control occasional invaders or aggregating pests as listed on this label.

**Application Rates for Outdoor Pests:**

	<b>Concentration of Active Ingredient</b>	<b>Dilution Rate in Water</b>	<b>Application Method(s)</b>
<b>Treating Exterior Walls, Foundations and Structures</b>	0.025 to 0.050%	0.5 to 1.0 fl. oz./gallon or 0.5 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *General Surface Spray *Crack and Crevice Treatment *Spot Treatment
<b>Treating Nests and Harborage Areas of Stinging Insects</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	
<b>To Control Swarming Termites and Wood Destroying Pests</b>	0.050%	1.0 fl. oz./gallon or 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *General Surface Spray *Crack and Crevice Treatment *Spot Treatment
<b>Treating Ornamental Trees and Landscapes</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *Broadcast Spray Treatment *Foliar Spray Treatment
<b>Mosquito Breeding Sites</b>	0.025%	0.5 fl. oz./gallon or 0.5 fl. oz. in sufficient water to cover 1,000 sq. ft.	Apply as a coarse, wet spray. *Broadcast Spray Treatment *Foliar Spray Treatment
<b>Treating Lawns and Turf</b>	0.005 to 0.050%	0.1 to 1.0 fl. oz./gallon or 0.1 to 1.0 fl. oz. in sufficient water to cover 1,000 sq. ft. 4.3 to 43 fl. oz. per acre	Apply as a coarse, wet spray. *Broadcast Spray Treatment

**Treating exterior walls, foundations and structures:**

To control infestations of listed pests, treat exterior surfaces of buildings, walls, window frames, around garbage cans, eaves, cracks and crevices, porches, decks, gazebos, patios, carports, garages, fence lines, and other areas where pests are active or may be hiding.

To help prevent insect infestation of buildings: Treat a 2 to 6 foot band of soil or other substrate adjacent to buildings. Treat building foundation to a height of 2 to 3 feet. Treat exterior walls, eaves, cracks and crevices, and other areas where listed pests are active, and may find entrance into building.

Apply with sufficient water to adequately cover the area being treated, but do not allow dripping or runoff to occur. Alternate mixing directions are to use 2 to 4 fluid ounces of concentrate per 50 gallons of water.

**Stinging Insect Control Outdoors:**

For stinging insects, apply spray solution to nests and harborage areas late in the evening when stinging insects are commonly at rest. Thoroughly spray nest, nest entrance, harborage sites, and surrounding areas where insects land or walk. For nests inside walls, inject sufficient spray to treat nest. Do not use in areas where an electrical shock hazard exists. For applications made when pests are active, applicator should wear protective equipment as required. Repeat application when there are signs of renewed insect activity.

For Yellow Jacket control, ONSLAUGHT® Microencapsulated Insecticide can be mixed with baits in traps. Follow trap instructions for preparation of bait.

For control of Fire Ants, combine broadcast application with mound drenches to control foraging workers and newly mated fly-in queens. Apply ONSLAUGHT® Microencapsulated Insecticide as a broadcast application at a rate of 8 fl. oz. of concentrate per 1,000 sq. ft. Treat mounds with a registered fire ant mound treatment.

**Insect Control in Ornamental Trees and Landscapes:**

For residential and commercial trees, shrubs, ground covers, and bedding and foliage plants that will not be harvested for food, apply ONSLAUGHT® Microencapsulated Insecticide in appropriate volumes of water to obtain thorough coverage. Apply as a full-coverage foliar spray, applying to the point of drip but not runoff. Treat active soil pests with an application to surrounding soil. Repeat treatments as necessary to achieve control, using higher application rates as pest pressure and foliage area increases. Repeat applications as necessary, but no more than once every seven (7) days. Certain plants may be sensitive to the final spray solution. A small-scale test is recommended to verify safety to ornamental plants. Spray and observe for one (1) week prior to application of an entire planting.

**Insect Control on Lawns, Turf Grass, and Turf:**

For best results, lawn or turf grass should be mowed 1-2 days before spraying. Treat with spraying equipment or a hand sprayer. Use application volumes of up to 10 gallons per 1,000 sq. ft. to get uniform coverage when treating dense grass foliage. For low volume applications using less than 2 gallons of spray solution per 1,000 sq. ft., immediate irrigation with at least 0.25 inch of water is recommended to improve effectiveness on sub-surface pests.

When hand spraying, spray using a slow, even sweeping motion, making sure to cover the entire lawn or turf grass area where pests are observed. Spray under ornamentals and trees. Repeat treatments may be necessary at 7 to 14 day intervals.

**Rate Range:**

Use lower rate range for pests that are commonly exposed and will be contacted by spray solution at the time of application. For pests that will not be contacted by spray solution at the time of application, use the upper rate range. Use 1.0 fl. oz. per 1,000 sq. ft. when treating Mole Crickets and Chinch Bugs.

**To Kill Swarming Termites: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray when swarming termites are seen emerging from woodpiles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new reproductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood. This treatment is not a substitute for a comprehensive termite control program.

**To Kill Wood Destroying Pests: OUTDOORS ONLY.**

Apply spray mix as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also spray around doors, window and door frames, and other areas where these pests may hide or enter the house. Spray into cracks and crevices and, if necessary, drill small holes and spray into inaccessible wooden structural voids where these pests are suspected.

**Mosquito Breeding sites:**

Mosquito populations may be reduced by application of ONSLAUGHT® Microencapsulated Insecticide to sites where mosquitoes rest, harbor, and breed. Apply spray solution into tall grass, shrubbery, and around backyards and lawns where these pests may hover or rest. Apply while air is still.

Not for wide area mosquito abatement using aerial or truck mounted cold aerosol ULV sprayers and thermal fogging devices.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry area under lock and key. Post as a pesticide storage area. Always store pesticides in the original container. Store away from food, pet food, feed, seed, fertilizers, and veterinary supplies. Place liquid formulations on lower shelves and dry formulations above.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local government or by industry).

**CONTAINER HANDLING**

Nonrefillable container. Do not reuse or refill this container.

Triple rinse as follows: Empty the remaining contents into the application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinse into the application equipment or a mix tank or store rinse for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances.



# Onslaught®

## MICROENCAPSULATED INSECTICIDE

■ **Kills German roaches for up to 12 months (1 year) on indoor, unsealed concrete and/or cement surfaces only.**

**ACTIVE INGREDIENT:**

(S)-cyano(3-phenoxyphenyl)methyl-(S)-4-chloro-alpha-(1-methylethyl)benzeneacetate ..... 6.40%

**\*OTHER INGREDIENTS** ..... 93.60%

100.00%

\*Contains petroleum distillates

## KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BOOKLET FOR FIRST AID STATEMENT AND DIRECTIONS FOR USE

EPA Reg. No. 1021-1815  
Rev. 0613-0214  
F28371-107

EPA Est. No. 1021-MN-2<sup>A</sup>, 63416-MN-1<sup>B</sup>, 70815-GA-1<sup>C</sup>,  
39578-TX-1<sup>D</sup>, 44616-MO-1<sup>E</sup>, 47000-IA-1<sup>F</sup>  
Superscript is the first character of the lot number.

### NET CONTENTS: 1.0 GALLON

Manufactured by:  
8810 Tenth Avenue North • Minneapolis, MN 55427

